

Universitat de Lleida

Document downloaded from:

<http://hdl.handle.net/10459.1/68340>

The final publication is available at:

<https://doi.org/10.1038/s41586-019-1128-0>

Copyright

(c) Steidinger, B.S. et al., under exclusive licence to Springer Nature Limited, 2019

Title: Climatic controls of decomposition drive the global biogeography of forest tree symbioses

Authors: Steidinger BS^{1*}, Crowther TW^{2,†*}, Liang J^{3,4,*}, Van Nuland ME¹, Werner GDA⁵, Reich PB^{6,7}, Nabuurs G⁸, de-Miguel S^{9,10}, Zhou M³, Picard N¹¹, Herault B¹², Zhao X⁴, Zhang C⁴, Routh D² Abegg, Meinrad; Adou Yao, C. Yves; Alberti, Giorgio; Almeyda Zambrano, Angelica; Alvarez-Davila, Esteban; Alvarez-Loayza, Patricia; Alves, Luciana F.; Ammer, Christian; Antón-Fernández, Clara; Araujo-Murakami, Alejandro; Arroyo, Luzmila; Avitabile, Valerio; Aymard, Gerardo; Baker, Timothy; Balazy, Radomir; Banki, Olaf; Barroso, Jorcely; Bastian, Meredith; Bastin, Jean-Francois; Birigazzi, Luca; Birnbaum, Philippe; Bitariho, Robert; Boeckx, Pascal; Bongers, Frans; Bouriaud, Olivier; Brancalion, Pedro H. S.; Brandl, Susanne; Brearley, Francis Q.; Brien, Roel; Broadbent, Eben; Bruelheide, Helge; Bussotti, Filippo; Cazzolla Gatti, Roberto; Cesar, Ricardo; Cesljar, Goran; Chazdon, Robin; Chen, Han Y. H.; Chisholm, Chelsea; Cienciala, Emil; Clark, Connie J.; Clark, David; Colletta, Gabriel; Condit, Richard; Coomes, David; Cornejo Valverde, Fernando; Corral-Rivas, Jose J.; Crim, Philip; Cumming, Jonathan; Dayanandan, Selvadurai; de Gasper, André L.; Decuyper, Mathieu; Derroire, Géraldine; DeVries, Ben; Djordjevic, Ilija; Iêda, Amaral; Dourdain, Aurélie; Obiang, Nestor Laurier Engone; Enquist, Brian; Eyre, Teresa; Fandohan, Adandé Belarmain; Fayle, Tom M.; Feldpausch, Ted R.; Finér, Leena; Fischer, Markus; Fletcher, Christine; Fridman, Jonas; Frizzera, Lorenzo; Gamarra, Javier G. P.; Gianelle, Damiano; Glick, Henry B.; Harris, David; Hector, Andrew; Hemp, Andreas; Hengeveld, Geerten; Herbohn, John; Herold, Martin; Hillers, Annika; Honorio Coronado, Eurídice N.; Huber, Markus; Hui, Cang; Cho, Hyunkook; Ibanez, Thomas; Jung, Ilbin; Imai, Nobuo; Jagodzinski, Andrzej M.; Jaroszewicz, Bogdan; Johannsen, Vivian; Joly, Carlos A.; Jucker, Tommaso; Karminov, Viktor; Kartawinata, Kuswata; Kearsley, Elizabeth; Kenfack, David; Kennard, Deborah; Kepfer-Rojas, Sebastian; Keppel, Gunnar; Khan, Mohammed Latif; Killeen, Timothy; Kim, Hyun Seok; Kitayama, Kanehiro; Köhl, Michael; Korjus, Henn; Kraxner, Florian; Laarmann, Diana; Lang, Mait; Lewis, Simon; Lu, Huicui; Lukina, Natalia; Maitner, Brian; Malhi, Yadvinder; Marcon, Eric; Marimon, Beatriz Schwantes; Marimon-Junior, Ben Hur; Marshall, Andrew Robert; Martin, Emanuel; Martynenko, Olga; Meave, Jorge A.; Melo-Cruz, Omar; Mendoza, Casimiro; Merow, Cory; Monteagudo Mendoza, Abel; Moreno, Vanessa; Mukul, Sharif A.; Mundhenk, Philip; Nava-Miranda, Maria G.; Neill, David; Neldner, Victor; Nevenic, Radovan; Ngugi, Michael; Niklaus, Pascal; Oleksyn, Jacek; Ontikov, Petr; Ortiz-Malavasi, Edgar; Pan, Yude; Paquette, Alain; Parada-Gutierrez, Alexander; Parfenova, Elena; Park, Minjee; Parren, Marc; Parthasarathy, Narayanaswamy; Peri, Pablo L.; Pfautsch, Sebastian; Phillips, Oliver; Piedade, Maria Teresa; Piotto, Daniel; Pitman, Nigel C. A.; Polo, Irina; Poorter, Lourens; Poulsen, Axel Dalberg; Poulsen, John R.; Pretzsch, Hans; Ramirez Arevalo, Freddy; Restrepo-Correa, Zorayda; Rodeghiero, Mirco; Rolim, Samir; Roopsind, Anand; Rovero, Francesco; Rutishauser, Ervan; Saikia, Purabi; Saner, Philippe; Schall, Peter; Schelhaas, Mart-Jan; Schepaschenko, Dmitry; Scherer-Lorenzen, Michael; Schmid, Bernhard; Schöngart, Jochen; Searle, Eric; Seben, Vladimír; Serra-Diaz, Josep M.; Salas-Eljatib, Christian; Sheil, Douglas; Shvidenko, Anatoly; Silva-Espejo, Javier; Silveira, Marcos; Singh, James; Sist, Plinio; Slik, Ferry; Sonké, Bonaventure; Souza, Alexandre F.; *Stereńczak, Krzysztof*; *Svenning, Jens*-Christian; Svoboda, Miroslav; Targhetta, Natalia; Tchebakova, Nadja; Steege, Hans ter; Thomas, Raquel; Tikhonova, Elena; Umunay, Peter; Usoltsev, Vladimir; Valladares, Fernando; van der Plas, Fons; Van Do, Tran; Vasquez Martinez, Rodolfo; Verbeeck, Hans; Viana, Helder; Vieira,

Simone; von Gadow, Klaus; Wang, Hua-Feng; Watson, James; Westerlund, Bertil; Wiser, Susan; Wittmann, Florian; Wortel, Virginia; Zagt, Roderick; Zawila-Niedzwiecki, Tomasz; Zhu, Zhi-Xin; Zo-Bi, Irie Casimir and Peay KG¹†

Affiliations:

¹ Department of Biology, Stanford University, Stanford CA USA

² Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland

³ Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN, USA

⁴ Research Center of Forest Management Engineering of State Forestry Administration, Beijing Forestry University, Beijing, China

⁵ Department of Zoology, University of Oxford, Oxford UK

⁶ Department of Forest Resources, University of Minnesota

⁷ Hawkesbury Institute for the Environment, Western Sydney University

⁸ Wageningen University and Research

⁹ Departament de Producció Vegetal i Ciència Forestal, Universitat de Lleida-Agrotecnio Center

¹⁰ Forest Science and Technology Centre of Catalonia (CTFC)

¹¹ Food and Agriculture Organization of the United Nations

¹² Cirad, INP-HB, Univ Montpellier, UPR Forêts et Sociétés

*These authors contributed equally to this work and share the first-author

†Corresponding authors: Email kpeay@stanford.edu; albeca.liang@gmail.com; tom.crowther@usys.ethz.ch

The identity of the dominant microbial symbionts in a forest determines the ability of trees to access limiting nutrients from atmospheric or soil pools^{1,2}, sequester carbon^{3,4} and withstand the impacts of climate change¹⁻⁷. Characterizing the global distribution of symbioses, and identifying the factors that control it, are thus integral to understanding present and future forest ecosystem functioning. Here we generate the first spatially explicit map of forest symbiotic status using a global database of 1.2 million forest inventory plots with over 28,000 tree species. Our analyses indicate that climatic variables, and in particular climatically-controlled variation in decomposition rate, are the primary drivers of the global distribution of major symbioses. We estimate that ectomycorrhizal (EM) trees, which represent only 2% of all plant species⁸, constitute approximately 60% of tree stems on Earth. EM symbiosis dominates forests where seasonally cold and dry

climates inhibit decomposition, and are the predominant symbiosis at high latitudes and elevation. In contrast, arbuscular mycorrhizal (AM) trees dominate aseasonally warm tropical forests and occur with EM trees in temperate biomes where seasonally warm-and-wet climates enhance decomposition. Continental transitions between AM and EM dominated forests occur relatively abruptly along climate driven decomposition gradients, which is likely caused by positive plant-microbe feedbacks. Symbiotic N-fixers, which are insensitive to climatic controls on decomposition compared with mycorrhizal fungi, are most abundant in arid biomes with alkaline soils and high maximum temperatures. The climatically driven global symbiosis gradient we document represents the first spatially-explicit, quantitative understanding of microbial symbioses at the global scale and demonstrates the critical role of microbial mutualisms in shaping the distribution of plant species.

Microbial symbionts strongly influence the functioning of forest ecosystems. They exploit inorganic, organic² and/or atmospheric forms of nutrients that enable plant growth¹, determine how trees respond to elevated CO₂⁶, regulate the respiratory activity of soil microbes^{3,9}, and affect plant species diversity by altering the strength of conspecific negative density dependence¹⁰. Despite growing recognition of the importance of root symbioses for forest functioning^{1,6,11} and the potential to integrate symbiotic status into Earth system models that predict functional changes to the terrestrial biosphere^{11,12}, we lack spatially-explicit, quantitative maps of the different root symbioses at the global scale. Generating these quantitative maps of tree symbiotic states would link the biogeography of functional traits of belowground microbial symbionts with their 1.5 trillion host trees¹³, spread across Earth's forests, woodlands, and savannas.


The dominant guilds of tree root symbionts, arbuscular mycorrhizal (AM) fungi, ectomycorrhizal (EM) fungi, ericoid mycorrhizal (ErM) fungi, and nitrogen (N)-fixing bacteria (N-fixer) are all based on the exchange of plant photosynthate for limiting macronutrients. The AM symbiosis is the oldest of the four, having evolved nearly 500 million years ago, with EM, ErM and N-fixer plant taxa having evolved multiple times from an AM basal state. Plants that form the AM symbiosis are markedly more diverse than the other symbiotic groups, comprising nearly 80% of all terrestrial plant species, and principally rely on AM fungi for enhancing mineral phosphorus (P) uptake¹⁴. EM fungi evolved more recently from saprotrophic ancestors, and as a result may be better than AM fungi at competing with free living soil microbes for resources³. As such, some EM fungal lineages are more capable of mobilizing organic sources of soil nutrients (particularly nitrogen) compared with AM fungi^{15,16}. Association with EM fungi, but not AM fungi, has been shown to allow trees to accelerate photosynthesis in response to increased atmospheric CO₂ when soil nitrogen (N) is limiting⁶ and to inhibit soil respiration by decomposer microbes^{3,9} (but see ¹⁷). Because increased plant photosynthesis and decreased soil respiration both reduce atmospheric CO₂ concentrations, the EM symbiosis is associated with buffering the Earth's climate against anthropogenic changes.

In contrast to mycorrhizal fungi, which extract nutrients from the soil, symbiotic N-fixers (Rhizobia and Actinobacteria) convert atmospheric N₂ to plant-usable forms. Symbiotic N-fixers are responsible for a large fraction of biological soil-N inputs, which can increase N-availability in forests where they are locally abundant¹⁸. Both N-fixing bacteria and EM fungi often demand more plant photosynthate than does the AM symbiosis^{14,19,20}. Because tree growth and reproduction are limited by access to inorganic, organic and atmospheric sources of N, the distribution of these root symbioses is likely to reflect both environmental conditions that

maximize the cost-benefit ratio of symbiotic exchange as well as physiological constraints on different symbionts.

In one of the earliest efforts to understand the functional biogeography of plant root symbioses, Sir David Read²¹ categorically classified biomes by their perceived dominant mycorrhizal type and hypothesized that seasonal climates favor hosts associating with EM fungi due to their ability to compete directly for organic N. By contrast, it has been proposed that sensitivity to low temperatures has prevented N-fixers from dominating outside the tropics, despite the potential for N-fixation to alleviate N-limitation in boreal forests^{20,22}. However, global scale tests of these proposed biogeographic patterns and their proposed climate drivers are lacking or inconclusive²³⁻²⁵ and we have no understanding of the regional variations in this proposed latitudinal trend. To address this research gap, we compiled the first global ground-sourced survey database to reveal numerical abundances of each symbiosis across the global forested biomes, rather than incidence (presence or absence, e.g.,²³⁻²⁵), which is essential for identifying the shapes and potential mechanisms underlying transitions in forest symbiotic state along climatic gradients^{26,27}.

We determined the abundance of tree symbioses using GFBi, an extension from the plot-based Global Forest Biodiversity (GFB²⁸) database, which contains over 1.2 million forest inventory plots of individual-based measurement records from which we derive abundance information for entire tree communities (Figure 1).



146

147 **Figure 1. A map of 1 by 1 degree grid cells where we analyzed the proportion of tree stems**
 148 **and basal area for different symbiotic guilds (above). Circles show the location of training**
 149 **data, colored by geographic origin, while black squares show the extent of model**
 150 **projections. Panels below the map show actual vs. predicted proportion of basal area for**
 151 **ectomycorrhizal (EM), arbuscular mycorrhizal (AM), and N-fixer trees by continent and**
 152 **subregion, and demonstrate globally consistent model performance.**

Using published literature on the evolutionary histories of mycorrhizal and N-fixer symbioses^{8,25,29-33}, we assigned plant species from the GFBi to one of 5 symbiotic guilds: AM, EM, ErM, N-fixer, and non- or weakly-mycorrhizal (NM). Most plants with symbioses derived from the AM state retain the genetic potential to associate with AM fungi¹⁴. Thus, consistent with other studies in this field²⁹, we assigned tree species to the AM-exclusive guild if they were not EM, ericoid mycorrhizal, non-mycorrhizal, or N-fixers. While there is some uncertainty in such assignments, direct investigation of mycorrhizal status when done supports this assumption³⁴. Because individual measurements of mycorrhizal colonization are not possible at this scale, our models represent potential symbiotic associations.

To identify the key factors structuring symbiotic distributions we assembled 70 global predictor layers: 19 climatic (annual, monthly, and quarterly temperature and precipitation variables), 14 soil chemical (total soil N density, microbial N, C:N ratios and soil P fractions, pH, cation exchange capacity), 5 soil physical (soil texture and bulk density), 26 vegetative indices (leaf area index, total stem density, enhanced vegetation index means and variances), and 5 topographic variables (elevation, hillshade) (Table S7). Because decomposition is the dominant process by which soil nutrients become available to plants, we generated 5 additional layers that estimate the climatic control of decomposition. We parameterized decomposition coefficients according to the Yasso07 model^{35,36} using the following equation:

$$k = \text{Exp}(0.095T_i - 0.00014 T_i^2) (1 - \text{Exp}[-1.21 P_i]), \quad (1)$$

where P_i and T_i are precipitation and mean temperature, either quarterly or annually, and the constants $0.0095 (= \beta_1)$, $0.00014 (= \beta_2)$, and $-1.21 (= \gamma)$ are parameters fit using a previous global study of leaf litter mass-loss³⁶. Although local decomposition rates can vary significantly based on litter quality or microbial community composition³⁷, climate is the primary control at the

global scale³⁶. Decomposition coefficients describe how fast different chemical pools of leaf litter lose mass over time relative to a parameter, α , that accounts for leaf-chemistry. Decomposition coefficients (k) with values of 0.5 and 2 indicate a halving and doubling of decomposition rates relative to α , respectively (Supplemental Materials).

Given the large set of possible environmental predictors, we used the random forest machine-learning algorithm to identify the best predictors of global symbiosis distributions. The random forest algorithm averages multiple regression trees, each of which uses a random subset of all the model variables to predict a response. These regression trees identify optimal values along a predictor-gradient to “split” the model response into different nodes (e.g., predictions could be “split” into nodes of 50 or 75% of EM basal area depending on whether mean annual temperature is $>$ or $<$ 20°C). We ranked the importance of each variable according to inc node purity, which measures the decrease in model error that occurs whenever the response is split on that variable (Figure 2ABC). We first determined the influence and relationship of all 75 predictor layers on forest symbiotic state and then optimized our models using a stepwise reduction in variables, from least- to most-important. Soil chemical, vegetative, and topographic variables were the first to be eliminated from our models in this way. In a subsequent model that included only layers of climate, decomposition, and certain soil physical and chemical information, we found that the 4 most important variables accounted for $>85\%$ of the explained variability. We plot the partial-fits of these four variables for each symbiotic guild (Figure 2ABC).




Figure 2. Partial plots of residual variation explained by the four most important predictors of the proportion of tree basal area belonging to the (A) ectomycorrhizal (EM), (B) arbuscular mycorrhizal (AM), and (C) N-fixer symbiotic guilds. Variables are listed in declining importance from left to right, as determined by inc node purity, with points colored with a red-green-blue gradient according to their position on the x-axis of the most important variable (left-most panels for each guild), allowing cross visualization between predictors. Each panel lists two measures of variable importance, inc node purity (used for sorting) and %IncMSE (see Supplemental Information for description). Decomposition rates in (A) and (B) are in units of leaf litter mass loss per quarter. The abundance of each symbiont type transitions sharply along climatic gradients, suggesting that sites near the threshold are particularly vulnerable to switching their dominant symbiont guild with climate changes.

The three most numerically abundant tree symbiotic guilds each have reliable environmental signatures, with the four most important predictors accounting for 81, 79, and 52% of the total variability in EM, AM, and N-fixer relative basal area, respectively. Models for

ErM and NM lack strong predictive power given the relative rarity of these symbiotic states amongst trees, although the raw data do identify some local abundance hotspots for ErM (Figure S1). As a result, we focus the remainder of results and discussion on the three major tree symbiotic states (EM, AM, N-fixer). Despite the fact that data from N. and S. America constitute 65% of the training data (at the 1 by 1 degree grid scale), our models accurately predict the proportional abundances of the three major symbioses across all major geographic regions (Figure 1). The high performance of our models, which is robust to both K-fold cross-validation and rarefying samples so that all continents are represented with equal depth (Figures S10-12), suggest that regional variations in climate (including indirect effects on decomposition) and soil pH (for N-fixers) are the primary factors influencing the relative dominance of each guild at the global scale (geographic origin only explained ~2-5% of the variability in residual relative abundance) (Figure 1BCD, Table S8).

Random forest models should not be projected across predictor gradients that fall outside the ranges of their training data (e.g., grid cells with higher mean annual temperatures than the maximum used to fit the models). To prevent the over-projecting of our models over pixels where we lacked training data, we subset a global grid of predictor layers depending on whether (1) the grid cell fell within the top 60% of land surface with respect to tree stem density¹³ and either (2) fell within the univariate distribution of all the predictor layers from our training data and/or (3) fell within an 8-dimensional hypervolume defined by the unique set of the 4-best predictors of the relative abundance of each guild (Figure 2, Supplemental Materials). We then projected our models across only those grid cells that met these criteria, which constitutes 46% of the global land surface and 88% of global tree stems (Figure 1; Figure S16). While model validation indicates that our projections are robust, additional ground truthing of predictions to

identify any discrepancies would be incredibly valuable. If such discrepancies exist they can help fine tune climate-symbiosis models, or identify areas where climate might favor invasion by symbioses that have not yet evolved or dispersed to a particular biogeographic region.

In contrast to a recent global analysis of root traits, which concluded that plant evolution has favored reduced dependence on mycorrhizal fungi³⁸, we find that trees associating with the relatively more C-demanding and recently-derived EM fungi^{14,19} represent the dominant tree-symbiosis. By taking the average proportion of EM trees, weighted by spatially-explicit global predictions for tree stem density¹³, we estimate that approximately 60% of trees on earth are EM, despite the fact that only 2% of plant species associate with EM fungi (vs. 80% associating with AM fungi)^{8,29}. Outside of the tropics, the estimate for EM relative abundance increases to approximately 80% of trees.

Turnover among the major symbiotic guilds results in a tri-modal latitudinal abundance gradient, with the proportion of EM trees increasing (and AM trees decreasing) with distance from the equator, while N-fixing trees reach peak abundance in the arid zone around 30 degrees (Figure 3A, Figure 4). These trends are driven by abrupt transitional regions along continental climatic gradients (Figure 2), which skew the distribution of symbioses among biomes (Figure 3A) and drive strong patterns across geographic and topographic features that influence climate. For example, moving north or south from the equator, the first transitional zone separates warm (aseasonal), AM-dominated, tropical broadleaf forests (>75% median basal area, vs. 8% for EM trees) from the rest of the EM-dominated world forest system (Figure 2AB; Figure 3A). It stretches longitudinally across 25 degrees N and S, just beyond the dry tropical broadleaf forests (with 25% EM tree basal area; Figure 3A), where average monthly temperature variation reaches 3-5°C (Figure 2AB).




Figure 3. (A) The median proportion of tree basal area per biome (arranged in order of increasing mean temperature of the warmest quarter) for ectomycorrhizal (EM), arbuscular mycorrhizal (AM), and N-fixer symbiotic guilds. EM trees dominate all extra-tropical biomes. **(B)** The dependency of decomposition coefficients (k , solid and dotted lines) on temperature and precipitation during the warmest quarter with respect to predicted dominance of mycorrhizal symbiosis. AM forests transition to EM forests abruptly in the region between $k=1$ and 2, which is consistent with positive feedback between climatic and biological controls of decomposition.

Moving further N or S, the second transitional climate zone separates regions where decomposition coefficients during the warmest quarter of the year are less than 2 (see Figure 3B for the associated temperature and precipitation ranges). In N. America and China, this transition zone spans longitudinally around 50 degrees N, separating the mixed AM / EM temperate forests from their neighboring EM dominated boreal forests (75 vs 100% EM tree basal area, respectively; Figure 3A). This transitional decomposition zone bypasses W. Europe, which has temperature seasonality $> 5^{\circ}\text{C}$, but lacks sufficiently wet summers to accelerate decomposition coefficients beyond values associated with mixed AM/EM forests. The latitudinal transitions in symbiotic state observed among biomes are mirrored by within-biome transitions along elevation

gradients. For example, in tropical Mexico, warm and wet quarter decomposition coefficients < 2 occur along the slopes of the Sierra Madre, where mixed AM-exclusive and N-fixer woodlands in arid climates transition to EM dominated tropical coniferous forests (75% basal area, Figure 3A, Figure 4ABC, Figure S17-19). The southern hemisphere, which lacks the landmass to support extensive boreal forests, experiences a similar latitudinal transition in decomposition rates along the ecotone separating its tropical and temperate biomes, around 28 degrees S.

The abrupt transitions that we detected between forest symbiotic states along environmental gradients suggest that positive feedbacks may exist between climatic and biological controls of decomposition^{11,36}. In contrast to AM fungi, some EM fungi can use oxidative enzymes to mineralize organic nutrients from leaf litter, converting nutrients to plant-usable forms before transferring them to their host trees^{2,5}. Relative to AM trees, the leaf litter of EM trees is also chemically more resistant to decomposition, with higher C:N ratios and higher concentrations of decomposition-inhibiting secondary compounds¹¹. Thus, EM leaf litter can exacerbate climatic barriers to decomposition, promoting conditions where EM fungi have superior nutrient-acquiring abilities to AM-fungi^{5,11}. Such positive-feedbacks are known to cause abrupt ecosystem transitions along smooth environmental gradients between woodlands and grasses: trees suppress fires, which promotes seedling recruitment, while grass fuels fires, which kill tree seedlings³⁹. Our study provides the first evidence that rapid transitions in tree community structure along climate gradients could also be governed by positive-feedbacks between symbiotic guilds and nutrient cycling; although other types of interactions, such as environmentally sensitive competition hierarchies among symbiotic guilds, could also lead to abrupt transitions without specifically invoking feedback effects. In either case, the existence of abrupt transitions suggests that trees and associated microbial symbionts in transitional regions

along decomposition gradients should be susceptible to drastic turnover in symbiotic state with future environmental changes⁴⁰.

To illustrate the sensitivity of global patterns of tree symbiosis to climate change, we use the climate relationships we developed for current climate to project potential changes due to climate change. Relative to our global predictions using the most recent climate data, model predictions using the projected climates for 2070 suggest the abundance of EM trees will decline by as much as 10% (using a relative concentration pathway of 8.5 W/m²; Figure S25). Due to their position along decomposition gradients relative to the abrupt shift from EM to AM forests (Figure 2AB), our models predict the largest declines in EM abundance will occur along the boreal-temperate ecotone, although declines in species abundances can lag decades, or even centuries or millennia, behind associated climatic changes⁴¹. The predicted decline in EM trees corroborates the results of common garden transfer and simulated warming experiments, which demonstrate that some important EM hosts will decline at the boreal-temperate ecotone in altered climates⁴²⁻⁴⁴. Because of the low tree diversity in boreal forests, tree species loss around transition zones may have major consequences for forest related economic activity⁴⁵.




Figure 4. Predicted global maps (left) and latitudinal gradients (right, with solid line indicating the median and colored ribbon spanning the range from the 5% and 95% quantiles) of the proportion of tree basal area for (A) ectomycorrhizal (EM), (B) arbuscular mycorrhizal (AM), and (C) N-fixer symbiotic guilds.

The change in dominant nutrient exchange symbioses along climate gradients highlights the interconnection between atmospheric and soil compartments of the biosphere. The transition from AM to EM dominance corresponds with a shift from P to N limitation of plant growth with increasing latitude⁴⁶⁻⁴⁸. Including published global projections of total soil N or P, microbial N, or soil P fractions (labile, occluded, organic, and apatite) did not increase the amount of variation explained by the model or alter the variables identified as most important, and thus were dropped from our analysis. However, this does not necessarily mean that soil nutrient availability is unimportant at the global scale, as the best-available global data likely do not adequately represent local nutrient availability^{49,50}. Rather, our finding that climatic controls of decomposition best predict the dominant mycorrhizal associations mechanistically links symbiont physiology with climatic controls of soil nutrient release from leaf litter. These findings are consistent with Read's hypothesis²¹ that slow decomposition at high latitudes favors EM fungi due to their increased capacity to liberate organic nutrients². Thus, while more experiments are necessary to understand the specific mechanism by which nutrient competition favors dominance of AM or EM symbioses²⁶, we propose that the latitudinal and elevational transitions from AM to EM dominated forests be called Read's Rule.

While our analyses focus on prediction at large spatial scales appropriate to the available data, our findings with respect to Read's Rule also provide insight into how soil factors structure the fine-scale distributions of tree symbioses within our grid cells. For example, while at a coarse scale we find that EM trees are relatively rare in many wet tropical forests, individual tropical sites in our raw data span the full range from 0 – 100 % EM basal area. In much of the wet tropics, these EM dominated sites exist as outliers within a matrix of predominantly AM trees. In an apparent exception that proves Read's Rule, in aseasonal warm neotropical climates, which

accelerate leaf-decomposition and promote regional AM dominance (Figure 3), EM dominated tree stands can develop in sites where poor soils and recalcitrant litter slow decomposition and N mineralization^{26,51}. Landscape-scale variation in the relative abundance of symbiotic states also changes along climate gradients, with variability highest in xeric and temperate biomes (Figure S2), suggesting that the potential of local nutrient variability to favor particular symbioses is contingent on climate.

Whereas EM trees are associated with ecosystems where plant growth is thought to be primarily N-limited, N-fixer trees are not. Our results highlight the global extent of the “N-cycling paradox,” wherein some metrics suggest that N-limitation is greater in the temperate zone⁴⁶⁻⁴⁸, yet N-fixing trees are relatively more common in the tropics^{20,52,53} (Figure 3A). We find that N-fixers, which we estimate represent 7% of all trees, dominate forests with annual max temperatures >35°C and alkaline soils (particularly in North America and Africa, Figure 2C). They have the highest relative abundance in xeric shrublands (24%), tropical savannas (21%), and dry broadleaf forest biomes (20%), but are nearly absent from boreal forests (<1%) (Figure 3A, Figure 4). The decline in N-fixer tree abundance we observed with increasing latitude is also associated with a previously documented latitudinal shift in the identity of N-fixing microbes, from facultative N-fixing rhizobial bacteria in tropical forests to obligate N-fixing actinorhizal bacteria in temperate forests⁵². Our data are not capable of fully disentangling the several hypotheses that have been proposed to reconcile the N-cycling paradox^{20,54}. However, our results are consistent with the model prediction²² and regional empirical evidence^{27,55,56} that N-fixing trees are particularly important in arid biomes. Based primarily on the observed positive, nonlinear association of N-fixer relative abundance with the mean temperature of the hottest

month (Figure 2C), our models predict a two-fold increase in N-fixer relative abundance when transitioning from humid to dry tropical forest biomes (Figure 3A).

Although soil microbes are a dominant component of forests, both in terms of diversity and ecosystem functioning^{5,6,11}, identifying global-scale microbial biogeographic patterns remains an ongoing research priority. Our analyses confirm that Read's Rule, which is one of the first proposed biogeographic rules specific to microbial symbioses, successfully describes global transitions between mycorrhizal guilds. More generally, climate driven turnover among the major plant-microbe symbioses represents a fundamental biological pattern in the Earth system, as forests transition from low-latitude arbuscular mycorrhizal, to N-fixer, to high-latitude ectomycorrhizal ecosystems. The predictions of our model (which we make available as a global raster layer) can now be used to represent these critical ecosystem variations in global biogeochemical models used to predict climate-biogeochemical feedbacks within and between trees, soils, and the atmosphere. Additionally, the layer containing the proportion abundance of N-fixing trees can be used to map potential symbiotic N-fixation, which links together atmospheric pools of C and N. Future work can extend our findings to incorporate multiple plant growth forms and non-forested biomes, where similar patterns likely exist, to generate a complete global perspective. Our predictive maps leverage the most comprehensive global forest dataset to generate the first quantitative global map of forest tree symbioses, demonstrating how nutritional mutualisms are coupled with the global distribution of plant communities.

Acknowledgments

This work is supported in part by the Key Project of National Key Research and Development Plan, China (2017YFC0504005); the new faculty start-up grant, Department of Forestry and Natural Resources, Purdue University; Dept. of Energy (DOE) Biological and Environmental

381 Research Program Early Career Research Grant DE-SC0016097; DOB Ecology, Plant-for-the-
382 Planet and the German Ministry for Economic Development and Cooperation; São Paulo
383 Research Foundation, #2014/14503-7; São Paulo Research Foundation (FAPESP), #2003/12595-
384 7; Proyecto FONACIT No. 1998003436 and UNELLEZ No. 23198105; EU, Sumforest –
385 REFORM, Risk Resilient Forest Management, FKZ: 2816ERA02S; U.S. National Science
386 Foundation Long-Term Ecological Research grant DEB-1234162, German Science Foundation
387 (DFG), KROOF Tree and stand-level growth reactions on drought in mixed versus pure forests
388 of Norway spruce and European beech, PR 292/12-1; Bavarian State Ministry for Food,
389 Agriculture and Forestry, W07 longterm yield experiments, 7831-26625-2017 and Project No
390 E33; The Deutsche Forschungsgemeinschaft (DFG) Priority Program 1374 Biodiversity
391 Exploratories; The International Tropical Timber Organization, ITTO-Project PD 53/00 Rev.3
392 (F); The State Forest Management Centre, Estonia, and the Environmental Investment Centre,
393 Estonia; Natural Sciences and Engineering Research Council of Canada Discover Grant Project
394 (RGPIN-2014-04181 and STPGP428641); European Structural Funds by FEDER 2014-2020
395 GY0006894; European Investment Funds by FEDER/COMPETE/POCI-Operational
396 Competitiveness and Internationalization Programme, under Project POCI-01-0145-FEDER-
397 006958 and National Funds by FCT - Portuguese Foundation for Science and Technology, under
398 the project UID/AGR/04033/2013; Vietnam National Foundation for Science and Technology
399 Development (NAFOSTED-106-NN.06-2016.10); German Research Foundation (DFG, FOR
400 1246); The project LIFE+ ForBioSensing PL Comprehensive monitoring of stand dynamics in
401 Bialowieza Forest co-funded by Life Plus (contract number LIFE13 ENV/PL/000048) and the
402 National Fund for Environmental Protection and Water Management in Poland (contract number
403 485/2014/WN10/OP-NM-LF/D); National Natural Scientific Foundation of China (31660055

404 and 31660074); The Polish State Forests National Forest Holding (2016); The Dutch Ministry of
405 Economic Affairs for funding the Dutch National Forest Inventory; The Grant 11-TE11-0100
406 from the U.S. National Space and Aeronautics Administration; the Tropical Ecology,
407 Assessment, and Monitoring (TEAM) / Conservation International project for funding the data
408 collection, and the National Institut Research Amazon (INPA); The Ministère des Forêts, de la
409 Faune et des Parcs du Québec (Canada); The Exploratory plots of FunDiveEUROPE received
410 funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant
411 agreement 265171; DBT, Govt. of India through the project ‘Mapping and quantitative
412 assessment of geographic distribution and population status of plant resources of Eastern
413 Himalayan region’ (sanction order No. BT/PR7928/NDB/52/9/2006 dated 29th September 2006);
414 The financial support from Natural Sciences and Engineering Research Council of Canada to S.
415 Dayanandan; Czech Science Foundation Standard Grant (16-09427S) and European Research
416 Council advanced grant (669609); RFBR #16-05-00496; The project implementation
417 Demonstration object on the transformation of declining spruce forests into ecologically more
418 stable multifunctional ecosystems, ITMS 26220220026, supported by the Research &
419 Development Operational Program funded by the ERDF; The Swedish NFI, Department of
420 Forest Resource Management, Swedish University of Agricultural Sciences SLU; The National
421 Research Foundation (NRF) of South Africa (89967 and 109244) and the South African
422 Research Chair Initiative; University Research Committee of the University of the South Pacific,
423 and New Colombo Plan Funding through the Department of Foreign Affairs and Trade of the
424 Australian government; The TEAM project in Uganda supported by the Moore foundation and
425 Buffett Foundation through Conservation International (CI) and Wildlife Conservation Society
426 (WCS); COBIMFO project funded by the Belgian Science Policy Office (Belspo), contract no.

427 SD/AR/01A; The German Federal Ministry of Education and Research (BMBF) under Grant
428 FKZ 01LL0908AD for the project “Land Use and Climate Change Interactions in the Vu Gia
429 Thu Bon River Basin, Central Vietnam” (LUCCI); Programme Tropenbos Côte d'Ivoire : projet
430 04/97-1111a du “Complément d’Inventaire de la Flore dans le Parc National de Tai”; The Danish
431 Council for Independent Research | Natural Sciences (TREECHANGE, grant 6108-00078B to
432 JCS) and VILLUM FONDEN (grant 16549); ERC Advanced Grant 291585 (“T-FORCES”) and
433 a Royal Society-Wolfson Research Merit Award; RAINFOR plots supported by the Gordon and
434 Betty Moore Foundation and the U.K. Natural Environment Research Council (NERC), notably
435 NERC Consortium Grants ‘AMAZONICA’ (NE/F005806/1), ‘TROBIT’ (NE/D005590/1), and
436 ‘BIO-RED’ (NE/N012542/1); Fundação de Amparo à Pesquisa e Inovação de Santa Catarina,
437 FAPESC (2016TR2524), Conselho Nacional de Desenvolvimento Científico e Tecnológico,
438 CNPq [312075/2013-8]; “Investissement d’Avenir” grant managed by Agence Nationale de la
439 Recherche (CEBA, ref. ANR- 10-LABX-25-01); CIFOR's Global Comparative Study on
440 REDD+ funded by the Norwegian Agency for Development Cooperation (Norad), the Australian
441 Department of Foreign Affairs and Trade (DFAT), the European Union (EU), the International
442 Climate Initiative (IKI) of the German Federal Ministry for the Environment, Nature
443 Conservation, Building and Nuclear Safety (BMUB), and the CGIAR Research Program on
444 Forests, Trees and Agroforestry (CRP-FTA), and donors to the CGIAR Fund; The Nature and
445 Biodiversity Conservation Union (NABU) under the project entitled “Biodiversity under Climate
446 Change: Community Based Conservation, Management and Development Concepts for the Wild
447 Coffee Forests”, funded by the German Federal Ministry for the Environment, Nature
448 Conservation and Nuclear Safety (BMU) through the International Climate Initiative (IKI); The
449 Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); The institutional

450 project “EXTEMIT - K”, no. CZ.02.1.01/0.0/0.0/15_003/0000433 financed by OP RDE; EC DG
451 VIII grant BZ-5041 (ECOSYN), NWO-WOTRO (W84-204), and GTZ; AfriTRON network
452 plots funded by the local communities and NERC, ERC, European Union, Royal Society and
453 Leverhume Trust; BOLFOR (Proyecto de Manejo Forestal Sostenible- Bolivia); The Global
454 Environment Research Fund F-071 and D-1006, and JSPS KAKENHI Grant Numbers
455 JP17K15289; The National Institute of Biology(Now Research Center for Biology), LIPI
456 (Indonesian Institute of Sciences), Indonesia IFBN project (contract 4000114425/15/NL/FF/gp)
457 funded by ESA; NSF grant DBI-1565046; Swiss National Science Foundation (SNSF No.
458 130720, 147092); Projects D/9170/07, D/018222/08, D/023225/09 and D/032548/10 funded by
459 the Spanish Agency for International Development Cooperation [Agencia Española de
460 Cooperación Internacional para el Desarrollo (AECID)] and Fundación Biodiversidad, in
461 cooperation with the Universidad Mayor de San Simón (UMSS), the FOMABO (Manejo
462 Forestal en las Tierras Tropicales de Bolivia) project and CIMAL (Compañía Industrial
463 Maderera Ltda.); The Agency for Economic and Environmental Development (DDEE) of the
464 north province of New Caledonia (the projects Ecofor & Cogefor, 2011-2016); Russian Science
465 Foundation (16-17-10284 “The accumulation of carbon in forest soils and forest succession
466 status”); Norwegian Ministry of Food and Agriculture; A grant from the Royal Society and the
467 Natural Environment Research Council (UK) to S.L.L.; The Spanish Agency for International
468 Development Cooperation [Agencia Española de Cooperación Internacional para el Desarrollo
469 (AECID)] and Fundación Biodiversidad, in cooperation with the governments of Syria and
470 Lebanon; COBIMFO Project, Federal Science Policy, Belgium; Consejo Nacional de Ciencia y
471 Tecnología, Mexico; Comisión Nacional Forestal, Mexico; BEF-China project (FOR 891)
472 funded by the German Research Foundation (DFG); WWF Russell Train Fellowship to P.M.U.

(Grant ST54); Wildlife Conservation Society DRC Program under CARPE Funding; Seoul National University Big Data Institute through the Data Science Research Project 2016, R&D Program for Forest Science Technology (Project No. 2013069C10-1719-AA03 & S111215L020110) funded by Korea Forest Service (Korea Forestry Promotion Institute); The European Union's Horizon 2020 research and innovation program within the framework of the MultiFUNGtionality Marie Skłodowska-Curie Individual Fellowship (IF-EF) under grant agreement 655815; Tropenbos International-Suriname; The Institute for World Forestry, University of Hamburg; REMBIOFOR Project "Remote sensing based assessment of woody biomass and carbon storage in forests" funded by The National Centre for Research and Development, Warsaw, Poland, under the BIOSTRATEG program (agreement no. BIOSTRATEG1/267755/4/NCBR/2015); The National Science Centre, Poland (Grant: 2011/02/A/NZ9/00108); Project "Environmental and genetic factors affecting productivity of forest ecosystems on forest and post-industrial habitats" (2011-2015; no. OR/2717/3/11); Project "Carbon balance of the major forest-forming tree species in Poland" (2007-2011; no. 1/07) funded by the General Directorate of State Forests, Warsaw, Poland; and the research professorship for "Ecosystem-based sustainable development" funded by Eberswalde University for Sustainable Development. GK was supported by an Alexander von Humboldt fellowship. GDAW was supported from a Newton International Fellowship from the Royal Society.

We thank the following agencies, initiatives, teams, and individuals for data collection and other technical support: the Global Forest Biodiversity Initiative (GFBI) for establishing the data standards and collaborative framework; United States Department of Agriculture, Forest Service, Forest Inventory and Analysis (FIA) Program; University of Alaska Fairbanks; The SODEFOR,

496 Ivory Coast; the Queensland Herbarium and past Queensland Government Forestry and Natural
 497 Resource Management Departments and staff for data collection for over seven decades. Ziaur
 498 Rahman Laskar, Salam Dilip, Bijit, Bironjoy and Samar; Badru Mugerwa and Emmanuel
 499 Akampurira, together with a team of field assistants (Valentine and Lawrence); all persons who
 500 made the Third Spanish Forest Inventory possible, especially the main coordinator, J. A.
 501 Villanueva (IFN3); Italian and Friuli Venezia Giulia Forest Services (Italy); Rafael Ávila and
 502 Sharon van Tuylen, Insituto Nacional de Bosques (INAB), Guatemala for facilitating
 503 Guatemalan data; The National Focal Center for Forest condition monitoring of Serbia (NFC),
 504 Institute of Forestry, Belgrade, Serbia; The Thünen Institute of Forest Ecosystems (Germany) for
 505 providing National Forest Inventory data; All TEAM data provided by the Tropical Ecology
 506 Assessment and Monitoring (TEAM) Network, a collaboration between Conservation
 507 International, the Missouri Botanical Garden, the Smithsonian Institution, and the Wildlife
 508 Conservation Society, and partially funded by these institutions, the Gordon and Betty Moore
 509 Foundation, and other donors, with thanks to all current and previous TEAM site manager and
 510 other collaborators that helped collecting data; The people of the Redi Doti, Pierrekondre and
 511 Cassipora village who were instrumental in assisting with the collection of data and sharing local
 512 knowledge of their forest; and the dedicated members of the field crew of Kabo 2012 census.
 513 Yadvinder Malhi's contribution was supported by an ERC Advanced Investigator Award GEM-
 514 TRAITS (321131).

515 **References**

- 516 1 Batterman, S. A. et al. Key role of symbiotic dinitrogen fixation in tropical forest
 517 secondary succession. *Nature* **502**, 224-227, doi:10.1038/nature12525 (2013).
- 518 2 Shah, F. et al. Ectomycorrhizal fungi decompose soil organic matter using oxidative
 519 mechanisms adapted from saprotrophic ancestors. *New Phytol* **209**, 1705-1719,
 520 doi:10.1111/nph.13722 (2016).

521 3 Averill, C., Turner, B. L. & Finzi, A. C. Mycorrhiza-mediated competition between
522 plants and decomposers drives soil carbon storage. *Nature* **505**, 543-+,
523 doi:10.1038/nature12901 (2014).

524 4 Clemmensen, K. E. et al. Roots and associated fungi drive long-term carbon
525 sequestration in boreal forest. *Science* **339**, 1615-1618, doi:10.1126/science.1231923
526 (2013).

527 5 Cheeke, T. E. et al. Dominant mycorrhizal association of trees alters carbon and nutrient
528 cycling by selecting for microbial groups with distinct enzyme function. *New Phytol.*
529 **214**, 432-442, doi:10.1111/nph.14343 (2017).

530 6 Terrer, C., Vicca, S., Hungate, B. A., Phillips, R. P. & Prentice, I. C. Mycorrhizal
531 association as a primary control of the CO₂ fertilization effect. *Science* **353**, 72-74,
532 doi:10.1126/science.aaf4610 (2016).

533 7 Johnson, D. J., Beaulieu, W. T., Bever, J. D. & Clay, K. Conspecific negative density
534 dependence and forest diversity. *Science* **336**, 904-907 (2012).

535 8 Brundrett, M. C. in *Biogeography of Mycorrhizal Symbiosis* 533-556 (Springer, 2017).

536 9 Averill, C. & Hawkes, C. V. Ectomycorrhizal fungi slow soil carbon cycling. *Ecol Lett*
537 **19**, 937-947, doi:10.1111/ele.12631 (2016).

538 10 Bennett, J. A. et al. Plant-soil feedbacks and mycorrhizal type influence temperate forest
539 population dynamics. *Science* **355**, 181-184 (2017).

540 11 Phillips, R. P., Brzostek, E. & Midgley, M. G. The mycorrhizal-associated nutrient
541 economy: a new framework for predicting carbon-nutrient couplings in temperate forests.
542 *New Phytol.* **199**, 41-51, doi:10.1111/nph.12221 (2013).

543 12 Oleson, K. W. et al. Technical description of version 4.0 of the Community Land Model
544 (CLM). (2010).

545 13 Crowther, T. W. et al. Mapping tree density at a global scale. *Nature* **525**, 201 (2015).

546 14 Heijden, M. G., Martin, F. M., Selosse, M. A. & Sanders, I. R. Mycorrhizal ecology and
547 evolution: the past, the present, and the future. *New Phytol.* **205**, 1406-1423 (2015).

548 15 Lindahl, B. D. & Tunlid, A. Ectomycorrhizal fungi—potential organic matter
549 decomposers, yet not saprotrophs. *New Phytol.* **205**, 1443-1447 (2015).

550 16 Liu, X. et al. Partitioning of soil phosphorus among arbuscular and ectomycorrhizal trees
551 in tropical and subtropical forests. *Ecol Lett* **21**, 713-723 (2018).

552 17 Zhu, K., McCormack, M. L., Lankau, R. A., Egan, J. F. & Wurzburger, N. Association of
553 ectomycorrhizal trees with high carbon-to-nitrogen ratio soils across temperate forests is
554 driven by smaller nitrogen not larger carbon stocks. *Journal of Ecology* **106**, 524-535
555 (2018).

556 18 Binkley, D., Sollins, P., Bell, R., Sachs, D. & Myrold, D. Biogeochemistry of adjacent
557 conifer and alder-conifer stands. *Ecology* **73**, 2022-2033 (1992).

558 19 Leake, J. et al. Networks of power and influence: the role of mycorrhizal mycelium in
559 controlling plant communities and agroecosystem functioning. *Canadian Journal of*
560 *Botany* **82**, 1016-1045 (2004).

561 20 Hedin, L. O., Brookshire, E. N. J., Menge, D. N. L. & Barron, A. R. in *Annual Review of*
562 *Ecology Evolution and Systematics* Vol. 40 Annual Review of Ecology Evolution and
563 *Systematics* 613-635 (Annual Reviews, 2009).

564 21 Read, D. J. Mycorrhizas in Ecosystems. *Experientia* **47**, 376-391, doi:Doi
565 10.1007/Bf01972080 (1991).

566 22 Houlton, B. Z., Wang, Y.-P., Vitousek, P. M. & Field, C. B. A unifying framework for
567 dinitrogen fixation in the terrestrial biosphere. *Nature* **454**, 327 (2008).

568 23 Bueno, C. G. et al. Plant mycorrhizal status, but not type, shifts with latitude and
569 elevation in Europe. *Global Ecology and Biogeography* **26**, 690-699 (2017).

570 24 Soudzilovskaia, N. A., Vaessen, S., van't Zelfde, M. & Raes, N. in *Biogeography of*
571 *Mycorrhizal Symbiosis* 223-235 (Springer, 2017).

572 25 Brundrett, M. C. & Tedersoo, L. Evolutionary history of mycorrhizal symbioses and
573 global host plant diversity. *New Phytol.* (2018).

574 26 Peay, K. G. The mutualistic niche: mycorrhizal symbiosis and community dynamics.
575 *Annual Review of Ecology, Evolution, and Systematics* **47**, 143-164 (2016).

576 27 Pellegrini, A. F., Staver, A. C., Hedin, L. O., Charles-Dominique, T. & Tourgee, A.
577 Aridity, not fire, favors nitrogen-fixing plants across tropical savanna and forest biomes.
578 *Ecology* **97**, 2177-2183 (2016).

579 28 Liang, J. et al. Positive biodiversity-productivity relationship predominant in global
580 forests. *Science* **354**, aaf8957 (2016).

581 29 Brundrett, M. C. Mycorrhizal associations and other means of nutrition of vascular
582 plants: understanding the global diversity of host plants by resolving conflicting
583 information and developing reliable means of diagnosis. *Plant and Soil* **320**, 37-77,
584 doi:10.1007/s11104-008-9877-9 (2009).

585 30 Werner, G. D., Cornwell, W. K., Cornelissen, J. H. & Kiers, E. T. Evolutionary signals of
586 symbiotic persistence in the legume–rhizobia mutualism. *Proceedings of the National*
587 *Academy of Sciences* **112**, 10262-10269 (2015).

588 31 Afkhami, M. E. et al. Symbioses with nitrogen-fixing bacteria: nodulation and
589 phylogenetic data across legume genera. *Ecology* **99**, 502-502 (2018).

590 32 Tedersoo, L. et al. Global database of plants with root-symbiotic nitrogen fixation: Nod
591 DB. *Journal of Vegetation Science* (2018).

592 33 Wang, B. & Qiu, Y.-L. Phylogenetic distribution and evolution of mycorrhizas in land
593 plants. *Mycorrhiza* **16**, 299-363 (2006).

594 34 McGuire, K. et al. Dual mycorrhizal colonization of forest-dominating tropical trees and
595 the mycorrhizal status of non-dominant tree and liana species. *Mycorrhiza* **18**, 217-222
596 (2008).

597 35 Palosuo, T., Liski, J., Trofymow, J. & Titus, B. Litter decomposition affected by climate
598 and litter quality—testing the Yasso model with litterbag data from the Canadian intersite
599 decomposition experiment. *Ecological Modelling* **189**, 183-198 (2005).

600 36 Tuomi, M. et al. Leaf litter decomposition—estimates of global variability based on
601 Yasso07 model. *Ecological Modelling* **220**, 3362-3371 (2009).

602 37 Bradford, M. A. et al. Climate fails to predict wood decomposition at regional scales.
603 *Nature Climate Change* **4**, 625 (2014).

604 38 Ma, Z. et al. Evolutionary history resolves global organization of root functional traits.
605 *Nature* (2018).

606 39 Staver, A. C., Archibald, S. & Levin, S. Tree cover in sub-Saharan Africa: rainfall and
607 fire constrain forest and savanna as alternative stable states. *Ecology* **92**, 1063-1072
608 (2011).

609 40 Scheffer, M., Carpenter, S., Foley, J. A., Folke, C. & Walker, B. Catastrophic shifts in
610 ecosystems. *Nature* **413**, 591 (2001).

- 41 Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**, 37 (2003).
- 42 Reich, P. B. & Oleksyn, J. Climate warming will reduce growth and survival of Scots pine except in the far north. *Ecol Lett* **11**, 588-597, doi:10.1111/j.1461-0248.2008.01172.x (2008).
- 43 Fernandez, C. W. & Kennedy, P. G. Revisiting the 'Gadgil effect': do interguild fungal interactions control carbon cycling in forest soils? *New Phytol.* **209**, 1382-1394, doi:10.1111/nph.13648 (2016).
- 44 Reich, P. B. et al. Geographic range predicts photosynthetic and growth response to warming in co-occurring tree species. *Nature Climate Change* **5**, 148 (2015).
- 45 Hanewinkel, M. & Peyron, J.-L. Tackling climate change—the contribution of scientific knowledge in forestry. *Ann Forest Sci* **71**, 113-115 (2014).
- 46 Vitousek, P. M. Litterfall, nutrient cycling, and nutrient limitation in tropical forests. *Ecology* **65**, 285-298 (1984).
- 47 McGroddy, M. E., Daufresne, T. & Hedin, L. O. Scaling of C: N: P stoichiometry in forests worldwide: Implications of terrestrial redfield-type ratios. *Ecology* **85**, 2390-2401 (2004).
- 48 Reich, P. B. & Oleksyn, J. Global patterns of plant leaf N and P in relation to temperature and latitude. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 11001-11006 (2004).
- 49 Vitousek, P. M. & Sanford Jr, R. L. Nutrient cycling in moist tropical forest. *Annu Rev Ecol Syst* **17**, 137-167 (1986).
- 50 Grunwald, S., Thompson, J. & Boettinger, J. Digital soil mapping and modeling at continental scales: Finding solutions for global issues. *Soil Science Society of America Journal* **75**, 1201-1213 (2011).
- 51 Corrales, A., Mangan, S. A., Turner, B. L. & Dalling, J. W. An ectomycorrhizal nitrogen economy facilitates monodominance in a neotropical forest. *Ecol Lett* **19**, 383-392, doi:10.1111/ele.12570 (2016).
- 52 Menge, D. N., Lichstein, J. W. & Ángeles-Pérez, G. Nitrogen fixation strategies can explain the latitudinal shift in nitrogen-fixing tree abundance. *Ecology* **95**, 2236-2245 (2014).
- 53 Ter Steege, H. et al. Continental-scale patterns of canopy tree composition and function across Amazonia. *Nature* **443**, 444 (2006).
- 54 Menge, D. N. et al. Why are nitrogen-fixing trees rare at higher compared to lower latitudes? *Ecology* **98**, 3127-3140 (2017).
- 55 Liao, W., Menge, D. N., Lichstein, J. W. & Ángeles-Pérez, G. Global climate change will increase the abundance of symbiotic nitrogen-fixing trees in much of North America. *Global Change Biol* (2017).
- 56 Gei, M. et al. Legume abundance along successional and rainfall gradients in Neotropical forests. *Nature ecology & evolution*, 1 (2018).